

Συμπληρωματικό υλικό

Αραμπατζή Α., Καρούσου Ρ., Κουκ Κ.Μ., Χανλίδου Ε., Λαναράς Θ., Κοκκίνη Σ. 2014. Οι ποικίλες οσμές των φυτών «μέντας» στα φυτογεωγραφικά διαμερίσματα της Ελλάδας. 36ο Επιστημονικό Συνέδριο της Ελληνικής Εταιρείας Βιολογικών Επιστημών, Ιωάννινα, 8-10 Μαΐου 2014.

Βιβλιογραφία

Adam K, Sivropoulou A, Kokkini S, Lanaras T, Arsenakis M (1998) Antifungal activities of *Origanum vulgare* subsp. *hirtum*, *Mentha spicata*, *Lavandula angustifolia*, and *Salvia fruticosa* essential oils against human pathogenic fungi. J Agric Food Chem 46: 1739- 1745

Alexopoulos A, Kimbaris AC, Plessas S, Mantzourani I, Theodoridou I, Stavropoulou E, Polissiou MG, Bezirtzoglou E (2011) Antibacterial activities of essential oils from eight Greek aromatic plants against clinical isolates of *Staphylococcus aureus*. Anaerobe 17: 399- 402

Cook CM, Lanaras T, Kokkini S (2007) Essential oils of two *Calamintha glandulosa* (Req.) Bentham chemotypes in a wild population from Zakynthos, Greece. J Essent Oil Res 19: 534-539

Couladis M, Tzakou O (2001) Essential oil of *Calamintha nepeta* subsp. *glandulosa* from Greece. J Essent Oil Res, 13: 11-12

Dimopoulos P, Raus TH, Bergmeier E, Constantinidis TH, Iatrou G, Kokkini S, Strid A, Tzanoudakis D (2013) Vascular plants of Greece: an annotated checklist. – Berlin: Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Athens: Hellenic Botanical Society. Englera 31:1-370

Frantzios G, Mirotsoy M, Hatzia Apostolou E, Kral J, Scouras ZG, Mavragani- Tspidou P (1997) Insecticidal and genotoxic activities of mint essential oils. J Agric Food Chem 45: 2690- 2694

Hanlidou E, Kokkini S, Bosabalidis AM, Bessiere JM (1991) Glandular trichomes and essential oil constituents of *Calamintha menthifolia* (Lamiaceae). Plant Syst Evol 177: 17- 26

Kadoglidou K, Lagopodi A, Karamanoli K, Vokou D, Bardas GA, Menexes G, Constantinidou HA (2011) Inhibitory and stimulatory effects of essential oils and individual monoterpenoids on growth and sporulation of four soil-borne fungal isolates of *Aspergillus terreus*, *Fusarium oxysporum*, *Penicillium expansum*, and *Verticillium dahlia*. Eur J Plant Pathol 130: 297- 309

Kanakis CD, Petrakis EA, Kimbaris AC, Pappas C, Tarantilis PA, Polissiou MG (2011) Classification of Greek *Mentha pulegium* L. (Pennyroyal) samples, according to geographical location by fourier transform infrared spectroscopy

Karousou R, Grammatikopoulos G, Lanaras T, Manetas Y, Kokkini S (1998) Effects of enhanced uv-b radiation on *Mentha spicata* essential oils. Phytochemistry 8: 2273- 2277

Karousou R, Hanlidou E, Lazari D (2012) Essential-oil diversity of three *Calamintha* species from Greece. Chem Biodivers 9: 1364- 1372

Karousou R, Kokkini S, Bessiere JM, Vokou D (1996) *Calamintha cretica* (Lamiaceae), a Cretan endemic: Distribution and essential oil composition. Nord J Bot 16: 247- 252

- Karousou R, Lanaras T, Kokkini S (1998) Piperitone Oxide-Rich essential oils from *Mentha longifolia* subsp. *petiolata* and *M. x villosa-nervata* grown wild on the Island of Crete (S Greece) *J Essent Oil Res* 10: 375- 379
- Kokkalou E, Stefanou E (1990) The Volatile Oil of *Calamintha nepeta* (L.) Savi subsp. *glandulosa* (Req.) P.W. Ball, Endemic to Greece. *Flavour Fragr J* 5: 23- 26
- Kokkini S, Hanlidou E, Karousou R, Lanaras T (2002) Variation of pulegone content in pennyroyal (*Mentha pulegium* L.) plants growing wild in Greece. *J Essent Oil Res* 14: 224- 227
- Kokkini S, Hanlidou E, Karousou R, Lanaras T (2004) Clinal variation of *Mentha pulegium* essential oils along the climatic gradient of Greece. *J Essent Oil Res* 16: 588- 593
- Kokkini S, Karousou R, Lanaras T (1996) Essential oils of spearmint (Carvone-rich) plants from the Island of Crete (Greece). *Biochem Syst Ecol* 23: 425- 430
- Kokkini S, Karousou R, Lanaras T (1997) Essential oils with 1,2-epoxy-p-menthane derivatives from *Mentha spicata* plants growing across the Island of Crete. *Bot Acta* 110: 184- 189
- Koliopoulos G, Pitarokili D, Kioulos E, Michaelakis A, Tzakou O (2010) Chemical composition and larvicidal evaluation of *Mentha*, *Salvia*, and *Melissa* essential oils against the West Nile virus mosquito *Culex pipiens*. *Parasitol Res* 107: 327- 335
- Ntalli NG, Ferrari F, Giannakou I, Menkissoglu- Spiroudi U (2010) Phytochemistry and Nematicidal Activity of the Essential Oils from 8 Greek Lamiaceae Aromatic Plants and 13 Terpene Components. *J Agric Food Chem* 58: 7856- 7863
- Sivropoulou A, Kokkini S, Lanaras T, Arsenakis M (1995) Antimicrobial activity of mint essential oils. *J Agric Food Chem* 43: 2384- 2388
- Souleles C, Argyriadou N (1990) The volatile constituents of *Calamintha grandiflora*. *Planta Med* 56: 234- 5
- Souleles C, Argyriadou N, Philianos S (1987) Constituents of the essential oil of *Calamintha nepeta*. *J Nat Prod* 50: 510- 522
- Vokou D, Liotiri S (1999) Stimulation of soil microbial activity by essential oils. *Chemoecology* 9: 41- 45
- Vokou D, Varelzidou S, Katinakis P (1993) Effects of aromatic plants on potato storage: sprout suppression and antimicrobial activity. *Agric Ecosyst Environ* 47: 223- 235
- Yadav RKP, Halley JM, Karamanoli K, Constantinidou HI, Vokou D (2004) Bacterial populations on the leaves of Mediterranean plants: quantitative features and testing of distribution models. *Environ Exp Bot* 52: 63- 77
- Yadav RKP, Karamanoli K, Vokou D (2005) Bacterial colonization of the phyllosphere of Mediterranean perennial species as Influenced by leaf structural and chemical features. *Microb Ecol* 50: 185- 196
- Yadav RKP, Papatheodorou EM, Karamanoli K, Constantinidou HI, Vokou D (2008) Abundance and diversity of the phyllosphere bacterial communities of Mediterranean perennial plants that differ in leaf chemistry. *Chemoecology* 18: 217- 226